

- Inform your caregiver that in the event of a seizure, you should be placed on your side until the seizure is over.

SPECIAL CARE

CRANIOTOMY - CRANIOENCEPHALIC TRAUMA

PHYSICAL ACTIVITY

- Do activities that do not require much effort: watch TV or read.
- Your physical condition may vary, this is normal, due to swelling.

WOUND CARE

- Keep the wound covered, preventing it from getting wet.
- 3 days after surgery, uncover the wound and wash it with shampoo or soap and water.

GO TO THE EMERGENCY ROOM IF YOU HAVE:

- Repetitive headache, vomiting or seizures .
- Behavior changes: drowsiness, apathy, incoherence.
- Signs of wound infection or if the wound becomes progressively swollen.

VENTRICULOPERITONEAL SHUNT IN CHILDREN

WOUND CARE

- Keep the wound covered, preventing it from getting wet.
- 3 days after surgery, uncover the wound and wash it with shampoo or soap and water.

GO TO THE EMERGENCY ROOM IF YOU HAVE:

- Progressive swelling of the wound or the top of the head (in children).

TRANS-PHENOIDAL SURGERY

PHYSICAL ACTIVITY

Start your daily activities gradually, avoiding movements that require much effort (avoid bending over).

WOUND CARE

- Avoid blowing your nose. Sneeze with your mouth wide open.
- Apply saline solution to decongest the nose, if needed.
- To relieve bruising or swelling in the eyes, use warm clothes at night.

GO TO THE EMERGENCY ROOM IF YOU HAVE:

- Liquid coming out of the nose or salty taste when swallowing.
- Loss of vision.
- Severe headache, inability to move the neck, or discomfort to the light.

**Para comentarios y sugerencias contacte al
Servicio de Integración y Atención al Usuario - SIAU
Ext. 4190 - 4191 - 4192
siau@fvl.org.co**

Síguenos en:



fvlcali



fvltv



@FVLCali



@fundacionvalledelili



Fundacion Valle del Ili

Av. Simón Bolívar Cra 98 # 18-49 PBX (57)(2) 331 90 90

www.valledelili.org

331 9090 ext. 7333



Certificado N° 015